

National Qualifications 2019

X707/77/02

# Biology Section 1 — Questions

TUESDAY, 30 APRIL 1:00 PM – 3:30 PM

Instructions for the completion of Section 1 are given on *page 02* of your question and answer booklet X707/77/01.

Record your answers on the answer grid on page 03 of your question and answer booklet.

Before leaving the examination room you must give your question and answer booklet to the Invigilator; if you do not, you may lose all the marks for this paper.





## SECTION 1 — 25 marks Attempt ALL questions

- 1. Primary cell lines have
  - A a limited number of cell divisions and are sourced from tumours
  - B a limited number of cell divisions and are sourced directly from normal animal tissue
  - C an indefinite number of cell divisions and are sourced from tumours
  - D an indefinite number of cell divisions and are sourced directly from normal animal tissue.
- 2. The proteome is larger than the number of genes in the genome of an organism because
  - A not all genes are expressed as proteins in a particular cell
  - B post translational modifications generate multiple RNAs from a single gene
  - C alternative splicing generates multiple RNAs from a single gene
  - D each mRNA molecule is translated by many ribosomes.
- 3. A neuron in a squid, running from the brain to the tentacles, is 0.9 m long. A 25 mm length of neuron is depolarised every 0.001 s.

The time taken for an impulse to travel the length of this neuron is

- A 0.000028 s
- B 0.000036 s
- C 0.028 s
- D 0.036 s

- 4. The list shows some events that occur in the cell membrane of a neuron during nerve transmission.
  - 1. Binding of neurotransmitter to neuron
  - 2. Closure of voltage-gated ion channels
  - 3. Opening of ligand-gated ion channels
  - 4. Opening of voltage-gated ion channels

Which events contribute to the depolarisation of the resting potential of a neuron?

- A 1 and 2 only
- B 2 and 3 only
- C 1, 2 and 3 only
- D 1, 3 and 4 only
- 5. Which row in the table describes the expected effects of inhibition of the Na/KATPase?

	Intracellular Na ion concentration	Intracellular K ion concentration	Membrane polarity
Α	increase	decrease	increase
В	decrease	increase	increase
С	increase	decrease	decrease
D	decrease	increase	decrease

- 6. A decrease in the activity of glucose symport proteins in the cells lining the small intestine could be caused by an increase in the
  - A glucose concentration inside the small intestine
  - B sodium ion concentration in the cells
  - C ATP concentration in the cells
  - D potassium ion concentration in the cells.

[Turn over

7. The surface area to volume ratio of a cell is an important factor affecting transport into cells.

surface area to volume ratio =  $\frac{\text{surface area}(\mu m^2)}{\text{volume}(\mu m^3)}$ 

The surface area to volume ratio of an *E. coli* cell is 4.5. A eukaryotic cell has a surface area of  $1809 \,\mu\text{m}^2$  and a volume of  $7235 \,\mu\text{m}^3$ .

Compared to E. coli, the surface area to volume ratio of the eukaryotic cell is approximately

- A 1.1 times larger
- B 1.1 times smaller
- C 18 times larger
- D 18 times smaller.
- 8. Alpha helices in proteins are stabilised by
  - A hydrogen bonds
  - B ionic bonds
  - C disulphide bridges
  - D hydrophobic interactions.
- 9. Which row in the table describes the effects of an increase in temperature on haemoglobin and oxygen delivery to cells?

	Affinity of haemoglobin for oxygen	Oxygen delivery to tissue
Α	increase	increase
В	increase	decrease
С	decrease	increase
D	decrease	decrease

**10.** The lipid bilayers of cell membranes contain a number of different phospholipids, which are present in different proportions and distributed unevenly across the extracellular and intracellular faces of the bilayer.

Figures 1 and 2 show the proportions and distribution of four phospholipids in the cell membrane of human red blood cells.





Figure 1 Proportions of four phospholipids in the cell membrane





The key can be used to identify each of the phospholipids.

1	more than 25 % of total lipid less than 25 % of total lipid	go to 2 go to 3
2	found mainly on the extracellular face found mainly on the intracellular face	A B
3	found mainly on the extracellular face	C

3 found mainly on the extracellular face found mainly on the intracellular face


Which letter represents sphingomyelin?

[Turn over

D

Questions 11 and 12 refer to the following information.

Short-term variations in the time interval between heart beats is termed *heart rate variability* (HRV). HRV is one measure that can be used to assess health. The box plots give information from one study about the influence of age and gender on HRV.



- 11. From the data shown, which of the statements about HRV is correct?
  - A There is a trend towards lower HRV in females
  - B The greatest error in measurements is for ages 25–34
  - C The mean HRV for females aged 35–44 is lower than 40 units
  - D Gender-related differences in median values are greater than age-related differences
- 12. Which row in the table describes features of this study?

	Design of experiment	Type of data collected
Α	simple	ranked
В	simple	continuous
С	multifactorial	ranked
D	multifactorial	continuous

**13.** There are approximately 40 species of birds of paradise in New Guinea, on islands nearby, and in areas of mainland Australia. They are thought to have evolved from a crow-like common ancestor that lived 20 million years ago.

The list describes processes that are likely to have contributed to the evolution of the different species.

- X The food availability on a certain part of one island favoured the survival of male and female individuals with slender curved bills.
- Y On one island with abundant food choices, females choose mates whose head feathers have elongated plumes.
- Z Some males and females of a species of crow-like mainland bird were blown by a freak storm to some of the islands.

Which row in the table matches processes of evolution with descriptions from the list?

	Pro	ocesses of evoluti	on	
	Genetic drift	Selection		
	Genetic drift	Natural	Sexual	
А	Х	Z	Y	
В	Z	Х	Y	
С	Z	Y	Х	
D	Y	Х	Z	

**14.** Which row in the table indicates factors that can all lead to a high rate of evolution?

		Factor	
	Selection pressure	Generation time	Gene transfer
A	high	long	asexual reproduction
В	low	short	horizontal
С	low	long	horizontal
D	high	short	sexual reproduction

[Turn over

- 15. Which of the following adaptations can be explained using the Red Queen hypothesis?
  - 1. Acceleration of cheetahs is related to the speed of the antelope they prey on
  - 2. Migration of insectivorous birds is influenced by the availability of insects
  - 3. Resistance of insect larvae to parasitic wasp infection is a response to these wasps
  - 4. Ornamental head plumes of some male birds are attractive to females
  - A 1 and 2 only
  - B 1 and 3 only
  - C 2 and 3 only
  - D 3 and 4 only
- **16.** A student used a colorimetric assay to measure the activity of an enzyme in bananas. An extract was prepared from one banana and used five times to measure the enzyme activity. Three absorbance readings were taken for each of the five assays.

	Abs	orbance reading at 540	nm
Assay	1	2	3
1	0.55	0∙56	0.55
2	0.49	0.48	0.50
3	0.56	0.57	0.57
4	0.62	0.63	0.62
5	0.58	0.58	0.59

The data collected are shown in the table.

The student evaluated the data to be reliable and accurate.

The student's evaluation of the data is

- A correct because the data are reliable and accurate
- B incorrect because the data are accurate but not reliable
- C incorrect because the data are reliable but not necessarily accurate
- D incorrect because the data are not reliable and not necessarily accurate.

17. Biological fieldwork can sometimes be more dangerous than laboratory work.

Which of the following would **not** generally be involved in a risk assessment for carrying out fieldwork safely?

- A Identify hazards
- B Specify control measures
- C Assess safety training records of participants
- D Consider hazard severity and likelihood of occurring
- **18.** Some populations of a species may evolve to become more r-selected or K-selected depending on the nature of the habitat they occupy.


Which row in the table shows changes in the factors likely to be associated with a population becoming more K-selected?

	Environmental stability	Number of offspring
Α	increase	decrease
В	increase	increase
С	decrease	increase
D	decrease	decrease

[Turn over

**19.** The population of the snail *Vertigo antivertigo* was investigated at a small site in Wales. Equal numbers of quadrat samples were taken in three areas with different dominant vegetation and the number of individual snails in each was recorded.

The results are shown in the chart.



dominant vegetation in area

The information in the chart indicates that this species of snail prefers

- A short grass to long grass
- B irises and rushes to long grass
- C long grass to irises and rushes
- D short grass to irises and rushes.
- **20.** Biological fieldwork often requires the estimation of population size for a prey species. One method used is mark and recapture. If the method of marking reduced the camouflage coloration of this species, what effect would this be likely to have on the population estimate obtained?
  - A Recapture numbers would be increased and population size would be overestimated
  - B Recapture numbers would be increased and population size would be underestimated
  - C Recapture numbers would be decreased and population size would be overestimated
  - D Recapture numbers would be decreased and population size would be underestimated

21. In three-spined stickleback fish, males have a distinctive red underside in the breeding season, which is not present in females. Territorial males were presented with model fish, some of which had their undersides painted red and some of which were left unpainted. Males showed an automatic attack response to only the red-painted models.

This attack response behaviour is triggered by


- A sexual dimorphism
- B sexual selection
- C an honest signal
- D a sign stimulus.
- 22. The list shows three events in meiosis I.
  - Q Independent assortment
  - R Recombination of alleles of linked genes
  - S Pairing of homologous chromosomes

The order in which these events occur is

- A = S R Q
- B = S Q R
- C Q R S
- D Q S R.

[Turn over

**23.** The graph shows species richness of mollusc populations in areas of fenland. The number of species of mollusc was recorded in several areas as well as the calcium concentration in the fenland water. At each site the concentration of iron was also measured.



low iron concentration
high iron concentration


Which of the following generalisations can be drawn from the graph?

- A An increase in calcium from 100 to 400 mg/l increases the species richness
- B High iron concentration leads to the highest species richness
- C When both calcium and iron concentrations are high the species richness is highest
- D An increase in calcium up to 150 mg/l increases species richness
- 24. New patterns of resistance in *Plasmodium* have increased the challenge experienced in the treatment of malaria.

Which of the following strategies is **least** likely to reduce the challenges in the treatment and control of malaria?

- A Improved sanitation
- B Development of new culture methods for *Plasmodium*
- C Coordinated vector control
- D Building of new low-density housing in malarial areas

25. Which diagram shows the sequence of events in the scientific cycle?



#### [END OF SECTION 1. NOW ATTEMPT THE QUESTIONS IN SECTION 2 OF YOUR QUESTION AND ANSWER BOOKLET.]

-						
	FOR OFFICIAL USE	1				
	National Qualificatio 2019	ons			Mark	
X707/77/01			Sect	ion 1 —	Bi - Answe and Sec	iology er grid tion 2
TUESDAY, 30 APRIL						
1:00 PM - 3:30 PM						701*
						_
Fill in these boxes and rea	d what is printed I	below.	Town			
Forename(s)	Surnar	me			Number o	of seat
Date of birth						
Day Month	Year	Scottish ca	ndidate	e number		
Total marks — 90						
SECTION 1 —25 marks						

Attempt ALL questions. Instructions for the completion of Section 1 are given on *page 02*.

#### SECTION 2 — 65 marks

Attempt ALL questions.

A supplementary sheet for question 1 is enclosed inside the front cover of this question paper.

Write your answers clearly in the spaces provided in this booklet. Additional space for answers and rough work is provided at the end of this booklet. If you use this space you must clearly identify the question number you are attempting. Any rough work must be written in this booklet. Score through your rough work when you have written your final copy.

Use **blue** or **black** ink.

Before leaving the examination room you must give this booklet to the Invigilator; if you do not, you may lose all the marks for this paper.





The questions for Section 1 are contained in the question paper X707/77/02.

Read these and record your answers on the answer grid on page 03 opposite.

Use **blue** or **black** ink. Do NOT use gel pens or pencil.


- 1. The answer to each question is **either** A, B, C or D. Decide what your answer is, then fill in the appropriate bubble (see sample question below).
- 2. There is only one correct answer to each question.
- 3. Any rough working should be done on the additional space for answers and rough work at the end of this booklet.

#### Sample question

The thigh bone is called the


- A humerus
- B femur
- C tibia
- D fibula.

The correct answer is **B** — femur. The answer **B** bubble has been clearly filled in (see below).



#### Changing an answer

If you decide to change your answer, cancel your first answer by putting a cross through it (see below) and fill in the answer you want. The answer below has been changed to **D**.



If you then decide to change back to an answer you have already scored out, put a tick ( $\checkmark$ ) to the **right** of the answer you want, as shown below:









	Α	В	С	D
1	0	0	0	0
2	0	0	0	0
3	0	0	0	0
4	0	0	0	0
5	0	0	0	0
6	0	0	0	0
7	0	0	0	0
8	0	0	$\bigcirc$	0
9	0	0	0	0
10	0	0	0	0
11	0	0	0	0
12	0	0	0	0
13	0	0	0	0
14	0	0	0	0
15	0	$\bigcirc$	$\bigcirc$	0
16	0	0	0	0
17	0	$\bigcirc$	$\bigcirc$	0
18	0	0	0	0
19	0	$\bigcirc$	$\bigcirc$	0
20	0	0	0	0
21	0	0	0	0
22	0	0	0	0
23	0	0	0	$\bigcirc$
24	0	0	0	0
25	0	0	$\bigcirc$	0



DO NOT WRITE ON THIS PAGE



[Turn over for next question




				MARKS	WRITEIN			
			SECTION 2 — 65 marks		THIS MARGIN			
			Attempt ALL questions					
			Question 10 contains a choice					
•	Read through the supplementary sheet for question 1 before attempting this question.							
	(a)	Refe	r to Figure 2.					
			in a species there is usually a positive correlation between size and where larger organisms are older.					
		Use I	Figure 2 to explain if the data support this for veined squid.	1				
				-				
				-				
	(b)		researchers used evidence published by other scientists to support work on ageing squid.	:				
		the term used to describe published summaries of current knowledge recent findings in a particular field.	2 1					
	(c)	(i)	Explain why many marine organisms use external fertilisation.	1				
		(ii)	Give one cost of this type of fertilisation.	- 1				

\* X 7


07770106\*

1. (co	ontinue	d)	MARKS
(D)		r <b>to Figure 3.</b> Explain how <b>Figure 3a</b> suggests that larger males may have a reproductive advantage.	۱ 1
	(ii)	Does the data in <b>Figure 3b</b> also support this conclusion? Justify your answer.	- 1
(e)	Refei	r to Figure 4.	-
	(i)	Identify the month that would have the highest breeding activity.	1
	(ii)	Squid eggs can take 30 days to hatch. Explain how the data for April support this finding.	1
			-
	(iii)	The squid take over a month to mature from stage 2 to stage 3. Suggest a reason for the unexpected decrease in the proportions of squid at stage 2 caught in May.	f 1
		[Turn over	

Γ




(co	ntinued)	
(b)	If the procedure was not carried out correctly a positive result could occur in the absence of <i>F. hepatica</i> antigens; a <i>false positive</i> result.	
	Suggest a possible cause for this false positive result.	1
(c)	A pH buffer was used in all reagents and wash solutions. Explain why it is important to control pH in immunoassays.	2
		L
(a)	Infection with <i>F. hepatica</i> in cattle results in weight loss and a reduction in milk yield. Suggest a reason for the reduced milk yield in infected cattle.	1
	[Turn over	





# MARKS DO NOT WRITE IN THIS MARGIN (continued) 3. Figure 2 shows the depth of penetration in water of the wavelengths in daylight. Figure 2 white light (all colours) 300 350 400 450 500 550 600 650 700 0 100 depth (m) 200 300wavelength (nm) (c) Crab species tend to live in shallower coastal waters, whereas octopus species can live in open seas. 2 Explain how the data from Figures 1 and 2 support this statement. [Turn over





MARKS DO NOT THIS (continued) 4. (c) Cholera is a disease caused by the bacterium Vibrio cholerae. It causes severe watery diarrhoea, which can lead to dehydration and even death. The bacterium produces cholera toxin which interferes with the control of the CFTR protein channel by constantly activating a kinase enzyme. (i) Describe the reaction catalysed by a kinase enzyme. 1 (ii) Explain how the production of cholera toxin by Vibrio cholerae can lead to more water being drawn out of the epithelial cells. 2 [Turn over 07770113\* X 7






MARKS DO NOT WRITE IN THIS MARGIN Behaviour is an animal's response to internal and external stimuli. 6. Discuss how animal behaviour can be measured and recorded. 4 \* X 7 0 7 7 7 0 1 1 6 \*

[Turn over for next question



7. The figure shows the life cycle of the parasitic hairworm, *Paragordius varius*, found in lakes throughout North and South America.



- (a) Hairworm eggs develop into larvae that hatch and move slowly through water at the bottom of the lake, where they may be ingested by aquatic insect larvae. The insects act as intermediate hosts, transporting the parasites into the terrestrial environment when the aquatic insect larvae become flying adults. If the flying insects are eaten by crickets, the parasite larvae penetrate the gut of the cricket to enter the body cavity.
  - (i) Explain why the hairworm and the cricket can be described as symbionts.
  - (ii) Explain why the cricket, and not the aquatic insect larva, is the definitive host.



MARKS DO NOT

1

1

THIS

		MARKS
(con	tinued)	
	In the presence of water, the hairworm exits the cricket's body and commences the next stage of its life-cycle. Researchers have shown that the hairworm adults produce 'mind-controlling' chemicals, which cause their normally terrestrial cricket hosts to jump into bodies of water.	
	(i) Suggest how this modification of host behaviour benefits the parasite.	1
	(ii) Give another example of a host behaviour that can be altered as part of the extended phenotype of a parasite.	- - 1
	Adult hairworms are free-living in aquatic environments where they mate. Males die almost immediately after mating so only mate once. Females die shortly after laying eggs fertilised by a single male.	
	State the term used to describe this type of reproductive strategy.	1
	Hairworms are closely related to members of the phylum Nematoda. Name the phylum to which both their hosts belong.	1
	[Turn over	



8. The 1918 'Spanish flu' pandemic was caused by the influenza virus and is thought to have killed approximately 50 million people.



(a) (i) Humans have many defences that prevent viruses from entering the body including physical barriers and chemical secretions.

Name one of the non-specific immune responses that may result if the virus is able to bypass these defences.

(ii) Annual flu vaccinations are designed to give immunity to some strains of influenza for vulnerable individuals. The vaccines typically contain inactivated virus particles that have been purified to leave only two of the proteins that are found on the surface of the virus coat.

Suggest why a new flu vaccine needs to be developed annually.

2

1



MARKS DO NOT THIS 8. (continued) (b) Scientists have predicted that another flu pandemic is highly likely in the future. They are particularly concerned that this might arise from the H5N1 strain of influenza virus, which is common in wild bird species, and is able to pass from birds to humans. (i) What term is used to describe the level of harm caused to the host species by a virus? 1 (ii) The case fatality rate (CFR) is the proportion of infected individuals who die as a result of a disease. Spanish flu had an estimated CFR of 2.5% but the H5N1 flu strain has a CFR of 55%. If Spanish flu had been caused by the H5N1 strain, and it had infected a similar number of people, what would have been the resulting number of deaths? 1 Space for calculation [Turn over

\* X 7 0 7 7 7 0 1 2 1 \*

MARKS DO NOT THIS MARGIN 9. The sandperch, Parapercis cylindrica, is a fish that lives on the sea floor. Each female defends a territory in which it feeds and reproduces. Dependent on the population density of the species, a single male may defend a group of up to 10 neighbouring females. (a) All sandperch begin life as hermaphrodites and mature into females. What is meant by the term hermaphrodite? 1 (b) There is a strict size-based dominance hierarchy in the social groups; the mean size of males is greater than that of females. State the term used to describe this difference in mean size. 1 (c) If the male dies, the largest female will undergo a period of growth and will then change sex. State one **other** cause of sex change in organisms. 1 (d) A study was conducted to measure the growth of female fish undergoing sex-change. Social groups were created in a laboratory by placing groups of similar sized fish in tanks with identical environmental conditions. Two treatments were set up, each with 10 groups. Treatment 1 with 2 females and 1 male in each group Treatment 2 with 4 females and 1 male in each group On day 5, the male was removed from half of each of the treatment groups to create experimental groups. On day 35 the sex and growth of each fish was determined. The structure of the groups is shown in the table. Treatment 1 Treatment 2 (10 groups) (10 groups) Control Control Experimental Experimental (5 groups) (5 groups) (5 groups) (5 groups) 2 females Day 2 females 4 females 4 females 1 1 male 1 male 1 male 1 male Group Day 2 females 4 females 2 females 4 females structure 5 1 male 1 male Day 2 females 1 female 4 females 3 females



1 male

1 male


1 male

35

1 male

#### 9. (d) (continued)

At day 35 the growth of fish in each experimental group was compared to its control group. This was used to calculate a standardised value that took into account growth in the control group. The figure shows the mean standardised growth values for the experimental groups in treatments 1 and 2. Error bars represent standard error of the mean.



(i) Suggest why it was necessary to have separate control groups for treatments 1 and 2.

(ii) Give one conclusion that could be drawn about the effect of treatment 2 on the growth of sex-changing females compared to treatment 1.

(iii) Suggest an advantage of greater size in sex-changed females.



1

1

1

THIS

		MARK			
10.	Answer <b>either A or B</b> in the space below and on <i>page 25</i> .				
	Α	Discuss signalling between cells under the following headings.			
		(i) Hydrophilic signalling molecules and signal transduction <b>6</b>			
		(ii) Insulin signalling and diabetes 4			
	OR				
	В	Discuss control of the cell cycle under the following headings.			
		(i) Phases of the cell cycle and the importance of cell cycle checkpoints <b>6</b>			
		(ii) The role of cyclins and cyclin-dependent kinases 4			



### SPACE FOR ANSWER FOR QUESTION 10

[END OF QUESTION PAPER]



#### MARKS DO NOT WRITE IN THIS MARGIN

## ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORK



#### MARKS DO NOT WRITE IN THIS MARGIN

### ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORK



DO NOT WRITE ON THIS PAGE

Acknowledgement of copyright Question 8 Everett Historical/shutterstock.com

