FOR OFFICIAL USE

National
Qualifications
2019

X707/77/01

Biology

Section 1 — Answer grid and Section 2

TUESDAY, 30 APRIL 1:00 PM - 3:30 PM



Fill in these boxe	es and read w	hat is print	ed below.							
Full name of cent	tre			Town						
Forename(s)		Sui	rname				Nur	nber	of sea	at
Date of birth	1									
Day	Month	Year	Scottish ca	andidate	numbe	r				

Total marks — 90

SECTION 1 —25 marks

Attempt ALL questions.

Instructions for the completion of Section 1 are given on page 02.

SECTION 2 — 65 marks

Attempt ALL questions.

A supplementary sheet for question 1 is enclosed inside the front cover of this question paper.

Write your answers clearly in the spaces provided in this booklet. Additional space for answers and rough work is provided at the end of this booklet. If you use this space you must clearly identify the question number you are attempting. Any rough work must be written in this booklet. Score through your rough work when you have written your final copy.

Use blue or black ink.

Before leaving the examination room you must give this booklet to the Invigilator; if you do not, you may lose all the marks for this paper.





The questions for Section 1 are contained in the question paper X707/77/02.

Read these and record your answers on the answer grid on page 03 opposite.

Use blue or black ink. Do NOT use gel pens or pencil.


- 1. The answer to each question is **either** A, B, C or D. Decide what your answer is, then fill in the appropriate bubble (see sample question below).
- 2. There is **only one correct** answer to each question.
- 3. Any rough working should be done on the additional space for answers and rough work at the end of this booklet.

Sample question

The thigh bone is called the

- A humerus
- B femur
- C tibia
- D fibula.

The correct answer is **B** — femur. The answer **B** bubble has been clearly filled in (see below).



Changing an answer

If you decide to change your answer, cancel your first answer by putting a cross through it (see below) and fill in the answer you want. The answer below has been changed to **D**.



If you then decide to change back to an answer you have already scored out, put a tick (\checkmark) to the right of the answer you want, as shown below:





	Α	В	С	D
1	0	0	0	0
2	0	0	0	0
3	0	0	0	0
4	\circ	\circ	\circ	\circ
5				
6	0	0	0	0
7	0	0	0	0
8	0	0	0	0
9	0	0	0	0
10	0	0	0	\circ
11	0	0	0	0
12	0	0	0	0
13	0	0	0	0
14	0	0	\circ	\circ
15	0	0	0	0
16	0	0	0	0
17	0	0	0	0
18	0	0	0	0
19	0	0	0	0
20	0	0	0	0
21	0	0	0	0
22	0	0	0	\circ
23	0	0	0	0
24	0	0	0	0
25	0	0	0	0



[BLANK PAGE]

DO NOT WRITE ON THIS PAGE



[Turn over for next question

DO NOT WRITE ON THIS PAGE



SECTION 2 — 65 marks Attempt ALL questions Question 10 contains a choice

1.

	d thro stion.	ough the supplementary sheet for question 1 before attempting this	
(a)	Refe	r to Figure 2.	
		in a species there is usually a positive correlation between size and where larger organisms are older.	
	Use F	Figure 2 to explain if the data support this for veined squid.	1
(b)		researchers used evidence published by other scientists to support work on ageing squid.	
		the term used to describe published summaries of current knowledge recent findings in a particular field.	1
(c)	(i)	Explain why many marine organisms use external fertilisation.	1
	(ii)	Give one cost of this type of fertilisation.	1



	_			
4	(co	 .		1
1	 \boldsymbol{c}	ntii	וור	0 71
			II U	Cu


(d)	Refe	r to Figure 3.
	(i)	Explain how Figure 3a suggests that larger males may have a reproductive advantage.
	(;;)	Does the data in Figure 3b also support this conclusion?
	(11)	Justify your answer.
(e)	Refe	to Figure 4.
	(i)	Identify the month that would have the highest breeding activity.
	(ii)	Squid eggs can take 30 days to hatch. Explain how the data for April support this finding.
	(iii)	The squid take over a month to mature from stage 2 to stage 3. Suggest a reason for the unexpected decrease in the proportions of squid at stage 2 caught in May.
		Suggest a reason for the unexpected decrease in the proportions of



Fasciolosis is a disease of cattle caused by the flatworm Fasciola hepatica (F. hepatica).


F. hepatica antigens are found both in the blood and the milk of infected cattle and their presence is the basis of an enzyme-linked immunoassay used to identify infected animals. A positive assay is described in the figure.

1.




monoclonal antibodies specific to *F. hepatica* are bound to the assay plate

2.




an infected milk sample is added to the plate


3.



the plate is washed with a buffer



a second monoclonal antibody, specific to F. hepatica and linked to an enzyme, is added 5.



the plate is washed with a buffer

6.



the colourless enzyme substrate is added and is converted to a coloured product

(a) (i) The antibody added at step four is a monoclonal antibody.

State the meaning of the term monoclonal.

1

(ii) Monoclonal antibodies can be produced in a laboratory using hybridomas that are prepared by fusing together B lymphocytes and myeloma cells.

Name the chemical used to fuse these two cell types together.

1

2. ((continu	ed)
~ • '	(COLLCILLA	Cuj

occur in the absence of <i>F. hepatica</i> antigens; a <i>false positive</i> result.
Suggest a possible cause for this false positive result.
A pH buffer was used in all reagents and wash solutions.
Explain why it is important to control pH in immunoassays.
Infection with <i>F. hepatica</i> in cattle results in weight loss and a reduction in milk yield.
Suggest a reason for the reduced milk yield in infected cattle.



- 3. Photoreceptor system proteins are found across the three domains.
 - (a) (i) Name the light sensitive molecule in animals that combines with the protein opsin to form photoreceptors of the eye.

(ii) Rod cells contain rhodopsin.

Explain why these cells can function in low light intensity.


1

(iii) Describe the role of bacteriorhodopsin in archaea.

2

Figure 1 shows the absorption spectra of rhodopsin in a variety of organisms.

Figure 1




(b) Owlflies have rhodopsin that is sensitive to light of 345 nm (UV).
Suggest how the photoreceptor proteins of owlflies differ from those of crabs.

1

3. (continued)

Figure 2 shows the depth of penetration in water of the wavelengths in daylight.

Figure 2



(c) Crab species tend to live in shallower coastal waters, whereas octopus species can live in open seas.

Explain how the data from Figures 1 and 2 support this statement.

•	_		

[Turn over


2



1

- The CFTR gene encodes a large transmembrane protein (CFTR) with a symmetrical structure. This is composed of two transmembrane regions and two ATP-binding regions.
 - (a) Name the class of R-groups that predominate in the transmembrane regions of proteins.
 - (b) The protein CFTR is involved in the regulation of water content of extracellular mucus in the lungs and digestive system.

The figure represents the action of CFTR. It regulates the passage of chloride ions (Cl⁻) across membranes of epithelial (lining) cells. In order for this ion channel to open, the protein must bind two ATP molecules, as well as a phosphate group. The increased concentration of Cl⁻ outside the cell draws water out of the epithelial cells into the mucus, maintaining its fluidity.



- (i) What name is given to a molecule that binds to a protein?
- (ii) Suggest how the binding of ATP results in the opening of the chloride ion channel. 1

2

4. (c	ontinu	ed)
-------	--------	-----

Cholera is a disease caused by the bacterium Vibrio cholerae. It causes
severe watery diarrhoea, which can lead to dehydration and even death.
The bacterium produces cholera toxin which interferes with the control
of the CFTR protein channel by constantly activating a kinase enzyme.

(1)	Describe the reaction catalysed by a kinase enzyme.
, <u>.</u>	
(ii)	Explain how the production of cholera toxin by <i>Vibrio cholerae</i> can lead to more water being drawn out of the epithelial cells.




th re of	It is thought that the level of fruit and vegetable consumption could influence the risk of some diseases in humans. One large study has examined the relationship between the level of fruit and vegetable consumption and the risk of mortality. This was a <i>meta-analysis</i> , which is a type of observational study that combines and summarises data from several previous similar studies.						
(a	-	a-analyses provide good evidence and are widely used in emiological studies.					
	State	e what is meant by the term 'epidemiology'.	1				
fo in	or inclus icluded.	of previously published research papers was used to identify studies sion in the meta-analysis. However, not all possible studies were In most of the studies used, fruit and vegetable consumption was using questionnaires.					
(b		est why the use of questionnaires to collect information about fruit vegetable consumption may have led to some errors in the data.	1				
(c		possible criticism of meta-analyses is that the people carrying them select the previous studies to include, which could result in selection					
	(i)	Suggest one reason why an individual study might be considered invalid and therefore excluded from the meta-analysis.	1				
	(ii)	State the effect that selection bias would have on the sample obtained.	1				



1

5. (continued)

The results from this meta-analysis are shown in the graph. The risk of mortality was expressed as a hazard ratio, with a hazard ratio of 1.0 indicating no effect on the risk of mortality.



(d) The graph shows the 95% confidence intervals for the data.

The confidence interval for six servings per day is wider than that for two servings.

What does this indicate about these two sets of data?

(e) This study is consistent with the hypothesis that higher fruit and vegetable consumption is associated with a reduced risk of mortality over time.

Explain why this observational study can only suggest a possible link between fruit and vegetable consumption and mortality risk over time.

* X 7 0 7 7 7 0 1 1 5 *

		MARKS	WRITE I
6.	Behaviour is an animal's response to internal and external stimuli.		MARGIN
	Discuss how animal behaviour can be measured and recorded.	4	


[Turn over for next question

DO NOT WRITE ON THIS PAGE



1

7. The figure shows the life cycle of the parasitic hairworm, *Paragordius varius*, found in lakes throughout North and South America.



- (a) Hairworm eggs develop into larvae that hatch and move slowly through water at the bottom of the lake, where they may be ingested by aquatic insect larvae. The insects act as intermediate hosts, transporting the parasites into the terrestrial environment when the aquatic insect larvae become flying adults. If the flying insects are eaten by crickets, the parasite larvae penetrate the gut of the cricket to enter the body cavity.
 - (i) Explain why the hairworm and the cricket can be described as symbionts.

(ii) Explain why the cricket, and not the aquatic insect larva, is the definitive host.



7. (continued)

(b) In the presence of water, the hairworm exits the cricket's body and commences the next stage of its life-cycle. Researchers have shown that the hairworm adults produce 'mind-controlling' chemicals, which cause their normally terrestrial cricket hosts to jump into bodies of water.

(i) Suggest how this modification of host behaviour benefits the parasite.

1

(ii) Give another example of a host behaviour that can be altered as part of the extended phenotype of a parasite.

1

(c) Adult hairworms are free-living in aquatic environments where they mate. Males die almost immediately after mating so only mate once. Females die shortly after laying eggs fertilised by a single male.

State the term used to describe this type of reproductive strategy.

1

(d) Hairworms are closely related to members of the phylum Nematoda. Name the phylum to which both their hosts belong.

1



The 1918 'Spanish flu' pandemic was caused by the influenza virus and is thought to have killed approximately 50 million people.



(a) (i) Humans have many defences that prevent viruses from entering the body including physical barriers and chemical secretions.

> Name one of the non-specific immune responses that may result if the virus is able to bypass these defences.

1

(ii) Annual flu vaccinations are designed to give immunity to some strains of influenza for vulnerable individuals. The vaccines typically contain inactivated virus particles that have been purified to leave only two of the proteins that are found on the surface of the virus coat.

Suggest why a new flu vaccine needs to be developed annually.					
	Suggest why a	new flu vaccin	e needs to be	developed	annually

2



1

8. (continued)

- (b) Scientists have predicted that another flu pandemic is highly likely in the future. They are particularly concerned that this might arise from the *H5N1* strain of influenza virus, which is common in wild bird species, and is able to pass from birds to humans.
 - (i) What term is used to describe the level of harm caused to the host species by a virus?

(ii) The case fatality rate (CFR) is the proportion of infected individuals who die as a result of a disease. Spanish flu had an estimated CFR of 2.5% but the H5N1 flu strain has a CFR of 55%.

If Spanish flu had been caused by the *H5N1* strain, and it had infected a similar number of people, what would have been the resulting number of deaths?

Space for calculation



- The sandperch, Parapercis cylindrica, is a fish that lives on the sea floor. Each female defends a territory in which it feeds and reproduces. Dependent on the population density of the species, a single male may defend a group of up to 10 neighbouring females.
 - (a) All sandperch begin life as hermaphrodites and mature into females. What is meant by the term hermaphrodite?

- (b) There is a strict size-based dominance hierarchy in the social groups; the mean size of males is greater than that of females.
 - State the term used to describe this difference in mean size.

1

(c) If the male dies, the largest female will undergo a period of growth and will then change sex.

State one other cause of sex change in organisms.

1

(d) A study was conducted to measure the growth of female fish undergoing sex-change. Social groups were created in a laboratory by placing groups of similar sized fish in tanks with identical environmental conditions.

Two treatments were set up, each with 10 groups.

- Treatment 1 with 2 females and 1 male in each group
- Treatment 2 with 4 females and 1 male in each group

On day 5, the male was removed from half of each of the treatment groups to create experimental groups. On day 35 the sex and growth of each fish was determined. The structure of the groups is shown in the table.


		11.00	tment 1 groups)	Treatment 2 (10 groups)		
		Control (5 groups)	Experimental (5 groups)	Control (5 groups)	Experimental (5 groups)	
	Day 1	2 females 1 male	2 females 1 male	4 females 1 male	4 females 1 male	
Group structure	Day 5	2 females 1 male	2 females	4 females 1 male	4 females	
	Day 35	2 females 1 male	1 female 1 male	4 females 1 male	3 females 1 male	



1

9. (d) (continued)

At day 35 the growth of fish in each experimental group was compared to its control group. This was used to calculate a standardised value that took into account growth in the control group. The figure shows the mean standardised growth values for the experimental groups in treatments 1 and 2. Error bars represent standard error of the mean.



(i)	Suggest why	it was	necessary	to	have	separate	control	groups	foi
	treatments 1	and 2.							

(ii)	Give	one	conclusion	that	could	be	drawn	about	the	effect	of
	treat	ment	2 on the	growth	of se	x-ch	anging	females	cor	npared	to
	treat	ment	1.								

(iii)	Suggest an advantage of greater size in sex-changed females.	1



			MARKS	DO NOT WRITE IN THIS	
10.	Answer either A or B in the space below and on <i>page 25</i> .				
	A	Discuss signalling between cells under the following headings.			
		(i) Hydrophilic signalling molecules and signal transduction	6		
		(ii) Insulin signalling and diabetes	4		
	OR				
	В	Discuss control of the cell cycle under the following headings.			
		(i) Phases of the cell cycle and the importance of cell cycle checkpoints	6		
		(ii) The role of cyclins and cyclin-dependent kinases	4		

MARKS DO NOT WRITE IN THIS MARGIN SPACE FOR ANSWER FOR QUESTION 10 [END OF QUESTION PAPER]

MARKS DO NOT WRITE IN THIS MARGIN

ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORK



MARKS DO NOT WRITE IN THIS MARGIN ADDITIONAL SPACE FOR ANSWERS AND ROUGH WORK

[BLANK PAGE]

DO NOT WRITE ON THIS PAGE

Acknowledgement of copyright

Question 8 Everett Historical/shutterstock.com

